

Distance between trees (m)

Questionnaire for Field Sampling Prunings

Field data (Each questionnaire refers to one crop species cultivated per field)

Municipality :	Trento				
Latitude :	46.077530		<u></u>		
Longitude :	11.128635				
Parco Naturale Adamello Brenta Tre Riva del Garda Coogle Limone Sul Garda	to Ba	assano Grappa			
Field Size (ha)	0.95				
Crop Species	olives peache cherrie grapef	es 🔲	vineyards apricot oranges hazelnuts	apples nectarine tangerines chestnuts	pears plum lemons almonds
Variety of crop	-	Merlot			
Age of crop	-	20			
Density of crop (trees/ha)		3800			
Width between cultivated rows (m)					

Crop forms for vineyard Espalier Marquee Vase Crop forms for Olive Ancient olives Vase (1 stem) Vase (2-3 Bush stems from soil) (intensive 250-600 trees/ha) Superintensive (>1500 trees/ha) Crop forms for fruit trees Natural Vase Bush/Globe Spindle/Pyramid (very small trees)

Epsylon transversal

Palm/Fan

Soil Cover

Bare.No grass cover. Tillage several times per	Seasonal occurence. Herbicides+mowing <50%	>50% grass cover. Mowed several times per
year	soil cover	year
100% Grass cover. Mowed several times per		
vear		

	Crop	o Yield	
Average Crop yield (t/ha)			
Crop yield before measurement (t/ha) Ammount of product obtained for the year that the pruning measurement is perfomed in tonnes per hectare			
Irrigation	rain fed	partial irrigation	fully irrigated
Intensification degree Specify the amount of fertilizer and pesticides	organic high	low	intermediate
Pi	runing Operations Perfor	med (prior to measureme	ent)
Type of pruning	Maintenance Structuring Removal of old bran	Тор	afting oping oming
Pruning Method	Only manually	Mechanised pre- pruning + manual	Fully mechanised
Pruning Operations Specify the pruning operations that are carried out. Check as many as apply.	Manually shears pre-pruner:hedge trimmer	Assisted shears pre-pruner:discs	Chainsaw/ armchainsaw pre-pruning topping

Season of pruning Check as many as apply	January May September	February June October	March July November	April August December
Frequency of pruning	annual	biannual	biennial	Once per years
Was this type of pruning performed as usual?	Yes, it was performed as usual	No, less inte	ense No, mo than usual	re intense
Describe the ammount of prunings produced compared to other years	Much Less More	Less Much more	Same	
Reason for different ammount of prunings produced ?	Pruning Intensit Weather	у	Accumulation of pre	evious years

	Pruning Measurement
Date of Measurement (DD/MM/YY)	2011
Mode of measurement Specify the method used for pruning measurement	Per tree One or several single trees selected. Biomass per tree collected manually and weighted Per parcel (e.g. 100m2) in bags
	One or several parcels selected. Each parcel several trees. Biomass per parcel collected manually and weighted
	Per several rows (or in whole field) On large parcel, or a whole field is selected. Biomass collected and loaded to a truck. Weight of the load taken ia a scale for truck
Ammount of prunings obtained (t/ha) Specify the tonnes per hectare of prunin from the crop	2.11 Losses of prunings that weren't harvested (t/ha)
How many days the prunings were or measuring (days)	soil before
Moisture content (%) Specify the moisture content (%) of the scollected if available	omass

References-External links:Provide references on which the information is based on or highlight any comments

Raffaele Spinelli, Carla Nati, Luigi Pari, Enzo Mescalchin, Natascia Magagnotti, Production and quality of biomass fuels from mechanized collection and processing of vineyard pruning residues, Volume 89, Issue 1, January 2012, Pages 374–379 Data obtained from EuroPruning Biomass Ratios Database, through the Task 3.1 and Deliverable D3.1 Mapping and analysis of the pruning biomass potential in Europe.

