

Field data (Each questionnaire refers to one crop species cultivated per field)

Municipality : Koropi

Latitude : 37.8628

Longitude : 23.8652



Field Size (ha) 0.3

- Crop Species
- | | | | |
|--|------------------------------------|-------------------------------------|----------------------------------|
| <input checked="" type="checkbox"/> olives | <input type="checkbox"/> vineyards | <input type="checkbox"/> apples | <input type="checkbox"/> pears |
| <input type="checkbox"/> peaches | <input type="checkbox"/> apricot | <input type="checkbox"/> nectarine | <input type="checkbox"/> plum |
| <input type="checkbox"/> cherries | <input type="checkbox"/> oranges | <input type="checkbox"/> tangerines | <input type="checkbox"/> lemons |
| <input type="checkbox"/> grapefruit | <input type="checkbox"/> hazelnuts | <input type="checkbox"/> chestnuts | <input type="checkbox"/> almonds |

Variety of crop Megaritiki

Age of crop 60

Density of crop (trees/ha) 73

Width between cultivated rows (m) 7

Distance between trees (m) 9

Crop form

Crop forms for vineyard



Vase



Espalier



Marquee

Crop forms for Olive



Ancient olives



Vase (1 stem)



Vase (2-3 stems from soil)



Bush (intensive 250-600 trees/ha)



Superintensive (>1500 trees/ha)

Crop forms for fruit trees



Natural



Vase



Bush/Globe (very small trees)



Spindle/Pyramid



Palm/Fan



Epsilon transversal

Slope (%)

0

Soil Cover



Bare.No grass cover.
Tillage several times per
year



Seasonal occurrence.
Herbicides+mowing <50%
soil cover



>50% grass cover.
Mowed several times per
year



100% Grass cover.
Mowed several times per
year

Crop Yield

Average Crop yield (t/ha) 13.3

Crop yield before measurement (t/ha) 0

Amount of product obtained for the year that the pruning measurement is performed in tonnes per hectare

Irrigation rain fed partial irrigation fully irrigated

Intensification degree organic low intermediate
Specify the amount of fertilizer and pesticides high

Pruning Operations Performed (prior to measurement)

Type of pruning Maintenance Grafting
 Structuring Topping
 Removal of old branches Blooming

Pruning Method Only manually Mechanised pre-pruning + manual Fully mechanised

Pruning Operations
Specify the pruning operations that are carried out. Check as many as apply.



Manually shears



Assisted shears



Chainsaw/armchainsaw



pre-pruner:hedge trimmer



pre-pruner:discs



pre-pruning topping

Season of pruning
Check as many as apply

- | | | | |
|---|--|-----------------------------------|-----------------------------------|
| <input checked="" type="checkbox"/> January | <input checked="" type="checkbox"/> February | <input type="checkbox"/> March | <input type="checkbox"/> April |
| <input type="checkbox"/> May | <input type="checkbox"/> June | <input type="checkbox"/> July | <input type="checkbox"/> August |
| <input type="checkbox"/> September | <input type="checkbox"/> October | <input type="checkbox"/> November | <input type="checkbox"/> December |

Frequency of pruning

- | | | | |
|---------------------------------|-----------------------------------|-----------------------------------|---|
| <input type="checkbox"/> annual | <input type="checkbox"/> biannual | <input type="checkbox"/> biennial | <input checked="" type="checkbox"/> Once per 10 years |
|---------------------------------|-----------------------------------|-----------------------------------|---|

Was this type of pruning performed as usual ?

- | | | |
|--|--|--|
| <input checked="" type="checkbox"/> Yes, it was performed as usual | <input type="checkbox"/> No, less intense than usual | <input type="checkbox"/> No, more intense than usual |
|--|--|--|

Describe the amount of prunings produced compared to other years

- | | | |
|--|------------------------------------|-------------------------------|
| <input type="checkbox"/> Much Less | <input type="checkbox"/> Less | <input type="checkbox"/> Same |
| <input checked="" type="checkbox"/> More | <input type="checkbox"/> Much more | |

Reason for different amount of prunings produced ?

- | | |
|--|--|
| <input type="checkbox"/> Pruning Intensity | <input type="checkbox"/> Accumulation of previous years |
| <input type="checkbox"/> Weather | <input checked="" type="checkbox"/> age and development of the trees |

Pruning Measurement

Date of Measurement
(DD/MM/YY)

15/02/17

Mode of measurement

Specify the method used for pruning measurement

Per tree



One or several single trees selected. Biomass per tree collected manually and weighted

Number of trees : 4

Per parcel (e.g. 100m²) in bags



One or several parcels selected. Each parcel several trees. Biomass per parcel collected manually and weighted

Per several rows (or in whole field)



On large parcel, or a whole field is selected. Biomass collected and loaded to a truck. Weight of the load taken in a scale for trucks

Amount of prunings obtained (t/ha)

Specify the tonnes per hectare of prunings collected from the crop

4.27

Losses of prunings that weren't harvested (t/ha)

How many days the prunings were on soil before measuring (days)

21

Moisture content (%)

Specify the moisture content (%) of the biomass collected if available

Contact Data

Name : Georgakis Stamatis

Email : aoskoropiou@yahoo.gr

Phone : 6932832183

Profession : Koropi Agricultural Cooperative

Country : Greece

Olive production every two years. Pruning operations are performed the year after the olives are harvested.

It was performed removal of old branches pruning after 10 years. Larger branches (diameter > 6 cm) were collected before the measurement.

uP_running project field measurement performed by CERTH

